

THE DISCONTINUITY PROBLEM

VASCO BRATTKA 

Abstract. Matthias Schröder has asked the question whether there is a weakest discontinuous problem in the topological version of the Weihrauch lattice. Such a problem can be considered as the weakest unsolvable problem. We introduce the *discontinuity problem*, and we show that it is reducible exactly to the *effectively discontinuous problems*, defined in a suitable way. However, in which sense this answers Schröder's question sensitively depends on the axiomatic framework that is chosen, and it is a positive answer if we work in Zermelo–Fraenkel set theory with dependent choice and the axiom of determinacy AD. On the other hand, using the full axiom of choice, one can construct problems which are discontinuous, but not effectively so. Hence, the exact situation at the bottom of the Weihrauch lattice sensitively depends on the axiomatic setting that we choose. We prove our result using a variant of Wadge games for mathematical problems. While the existence of a winning strategy for Player II characterizes continuity of the problem (as already shown by Nobrega and Pauly), the existence of a winning strategy for Player I characterizes effective discontinuity of the problem. By *Weihrauch determinacy* we understand the condition that every problem is either continuous or effectively discontinuous. This notion of determinacy is a fairly strong notion, as it is not only implied by the axiom of determinacy AD, but it also implies Wadge determinacy. We close with a brief discussion of generalized notions of productivity.

§1. Introduction. The Weihrauch lattice has been used as a computability-theoretic framework to analyze the uniform computational content of mathematical problems from many different areas of mathematics, and it can also be seen as a uniform variant of reverse mathematics (a recent survey on Weihrauch complexity can be found in [9]). The notion of a mathematical problem has a very general definition in this approach.

DEFINITION 1.1 (Problems). A *problem* is a multi-valued function $f : \subseteq X \rightrightarrows Y$ on represented spaces X, Y that has a realizer.

We recall that by a *realizer* $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ of f , we mean a function F that satisfies $\delta_Y F(p) \in f \delta_X(p)$ for all $p \in \text{dom}(f \delta_X)$, where $\delta_X : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$ and $\delta_Y : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow Y$ are the representations of X and Y , respectively (i.e., partial surjective maps onto X and Y , respectively).

We note that we have added the condition here that a problem has to have a realizer, since we want to prove all our results over the base theory of Zermelo–Fraenkel set theory (ZF) together with the axiom of dependent choice (DC), if not otherwise

Received May 30, 2021.

2020 *Mathematics Subject Classification.* 03D30, 03D78, 03E15, 03E25, 03E60, 91A05.

Key words and phrases. Weihrauch complexity, axiom of determinacy, effective discontinuity, computable analysis, computability theory, games.

© The Author(s). 2022. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

0022-4812/23/8803-0011
DOI:10.1017/jsl.2021.106



mentioned. These axioms suffice to prove most results in Weihrauch complexity. Typically, the full axiom of choice (AC) is freely used in Weihrauch complexity, often just to guarantee the existence of some realizer. By $\langle p, q \rangle$ we denote the usual pairing function on $\mathbb{N}^{\mathbb{N}}$, defined by $\langle p, q \rangle(2n) = p(n)$, $\langle p, q \rangle(2n + 1) = q(n)$ for all $p, q \in \mathbb{N}^{\mathbb{N}}$, $n \in \mathbb{N}$. Weihrauch reducibility can now be defined as follows.

DEFINITION 1.2 (Weihrauch reducibility). Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq W \rightrightarrows Z$ be problems. Then f is called *Weihrauch reducible* to g , in symbols $f \leq_w g$, if there are computable $H, K : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $H \langle \text{id}, GK \rangle$ is a realizer of f whenever G is a realizer of g . Analogously, one says that f is *strongly Weihrauch reducible* to g , in symbols $f \leq_{sw} g$, if the expression $H \langle \text{id}, GK \rangle$ can be replaced by HGK . Both versions of the reducibility have topological counterparts, where one requires H, K only to be continuous and these reducibilities are denoted by \leq_w^* and \leq_{sw}^* , respectively.

The topological version of Weihrauch reducibility has always been studied alongside the computable version, and all four reducibilities induce a lattice structure (see [9] for references). Normally, the *Weihrauch lattice* refers to the lattice induced by \leq_w , but here we will freely use this term also for the lattice structure induced by \leq_w^* . If we want to be more precise, we will call the latter the *topological Weihrauch lattice*. Even though this lattice has been studied for about 30 years, very little is known about the structure of the lattice closer towards the bottom. Indeed Matthias Schröder has asked the following question¹ [6, Question 5.9].

QUESTION 1.3 (Matthias Schröder, 2018). *Does there exist a discontinuous problem f such that $f \leq_w^* g$ holds for any other discontinuous problem g ?*

Here a problem is called *continuous* if it has a continuous realizer and *discontinuous* otherwise. It is clear that the degree 0 of the nowhere defined problems f is the bottom degree of the (continuous) Weihrauch lattice. The second lowest degree, sometimes called 1, is the degree of the identity $\text{id} : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ that includes all somewhere defined continuous problems.² Essentially Schröder's question is whether there is a third degree 2 such that the topological Weihrauch lattice starts with the linear ordered structure $0 < 1 < 2$. We will prove that under certain conditions this is indeed so, namely there is such a third degree given by the problem

$$\text{DIS} : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}, p \mapsto \{q \in \mathbb{N}^{\mathbb{N}} : U(p) \neq q\}$$

that we call the *discontinuity problem* (see Definition 3.2). Here $U : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is a fixed universal computable function.

In Section 3 we prove that the discontinuity problem DIS characterizes effectively discontinuous problems in the following sense (see Theorem 3.4).

THEOREM 1.4 (Continuity and effective discontinuity). *Let $f : \subseteq X \rightrightarrows Y$ be a problem. Then we obtain*

¹The original question is phrased slightly differently, but we interpret it in the intended way.

²For the computability-theoretic and the strong topological version of Weihrauch reducibility the continuous problems with non-empty domain do not form a single equivalence class; however, together with 0 they still form the cone below id.




FIGURE 1. Problems $f : \subseteq X \rightrightarrows Y$ with respect to \leq_W^* and \leq_{sW}^* .

- (1) $f \leq_W^* id \iff f$ is continuous,
- (2) $DIS \leq_W^* f \iff f$ is effectively discontinuous.

The diagram in Figure 1 illustrates the situation. Here *effective discontinuity* of a problem f is defined in a very natural way (see Definition 3.1) using a continuous *discontinuity function* $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ that has to produce an input to every given potential continuous realizer of f on which this realizer fails.

In some sense the notion of effective discontinuity is reminiscent of the notion of *productivity* from classical computability theory, which can be regarded as the property of being “effectively not c.e.” Indeed a well-known theorem of Myhill [18] gives us the analog of Theorem 1.4 for subsets $A \subseteq \mathbb{N}$ and many-one reducibility \leq_m [25, Theorem 2.4.6], [29, Theorem 2.6.6]. Here the *halting problem* $K \subseteq \mathbb{N}$ plays the counterpart of the identity and $\mathbb{N} \setminus K$ the counterpart of DIS.

THEOREM 1.5 (Myhill, 1955). *Let $A \subseteq \mathbb{N}$. Then we obtain*

- (1) $A \leq_m K \iff A$ is c.e.,
- (2) $\mathbb{N} \setminus K \leq_m A \iff A$ is productive.

The diagram in Figure 2 illustrates the situation. In fact, Theorem 1.4 is proved with the help of the recursion theorem, in a similar way as Theorem 1.5. However, somewhat surprisingly, the recursion theorem is used for the implication “ \Leftarrow ” in the proof of Theorem 1.5(2) and for the direction “ \Rightarrow ” in the proof of Theorem 1.4(2).

It is well known that Theorem 1.5 does not express a dichotomy, i.e., there are sets which are neither c.e. nor productive. A set $A \subseteq \mathbb{N}$ is called *immune* if it is infinite, but does not include an infinite c.e. set [25, 29]. By a classical construction of Post immune sets exist [23], [25, Theorem 5.2.3], [29, Theorem 2.3.7] and they are clearly neither c.e. nor productive, since any productive set contains an infinite c.e. subset (see Figure 2).

Now a key question for us is whether there is a counterpart of immune sets in our situation, i.e., whether there are problems f that are discontinuous, but not effectively so (see the question mark in Figure 1). In fact, Post’s construction of a




FIGURE 2. Sets $A \subseteq \mathbb{N}$ with respect to many-one reducibility \leq_m .

simple set has some similarity to the construction of so-called *Bernstein sets* that can actually be used to construct discontinuous problems that are not effectively discontinuous (see Corollary 3.9). This leads to the following counterexample (see Theorem 3.10).

THEOREM 1.6. *Assuming ZFC there are problems $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ that are continuous, but not effectively so.*

In Section 4 we prove that the axiom of choice is actually required for such a construction. This can be achieved with the help of a variant of Wadge games for problems $f : \subseteq X \rightrightarrows Y$, originally considered by Nobrega and Pauly [20, 21]. In fact, we can prove the following result (see Theorem 4.3), part (1) of which is already due to Nobrega and Pauly.

THEOREM 1.7 (Wadge games). *We consider the Wadge game of a given problem $f : \subseteq X \rightrightarrows Y$. Then the following hold:*

- (1) f is continuous \iff Player II has a winning strategy for f ,
- (2) f is effectively discontinuous \iff Player I has a winning strategy for f .

This result implies that under the axiom of determinacy (AD), which states that every Gale–Stewart game is determined, i.e., either Player I or Player II has a winning strategy, we really obtain a dichotomy between continuity and effective discontinuity (see Corollary 4.11).

COROLLARY 1.8. *In $\text{ZF} + \text{DC} + \text{AD}$ every problem $f : \subseteq X \rightrightarrows Y$ is either continuous or effectively discontinuous.*

This can be proved by a suitable reduction of the Wadge game to a Gale–Stewart game. By *Weihrauch determinacy* we understand the condition that every problem $f : \subseteq X \rightrightarrows Y$ is either continuous or effectively discontinuous. Then Corollary 1.8 can also be rephrased such that AD implies Weihrauch determinacy. On the other hand, it is easy to see that Wadge games for problems $f : \subseteq X \rightrightarrows Y$ generalize Wadge games for subsets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ as originally considered by Wadge and hence




FIGURE 3. Determinacy properties in $ZF + DC$.

Weihrauch determinacy implies Wadge determinacy, which means that every Wadge game for subsets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ is determined (see Figure 3).

We leave it open how Weihrauch determinacy is exactly related to the other given notions of determinacy. We just mention that it is not known whether Wadge determinacy implies AD, and in fact there is the following conjecture attributed to Solovay (by Andretta [1]³).

CONJECTURE 1.9 (Solovay). In $ZF + V = L(\mathbb{R})$ Wadge determinacy is equivalent to the axiom of determinacy AD.

We have some partial results (see Proposition 4.15) that suggest that Weihrauch determinacy is actually even closer related to AD than Wadge determinacy.

In any case, our results show that the exact situation at the bottom of the topological Weihrauch lattice sensitively depends on the underlying axioms. Using AD we obtain the following result (as a consequence of Corollary 1.8).

THEOREM 1.10. In $ZF + DC + AD$ the topological Weihrauch lattice starts with three linearly ordered degrees, represented by $0 \leq_w^* \text{id} \leq_w^* \text{DIS}$.

If we replace AD by the axiom of choice AC, then the linear part of the order at the bottom is just $0 \leq_w^* \text{id}$ and the situation becomes more complicated afterwards (and by Theorem 3.10 this is even true if we move to the coarser parallelized version of the Weihrauch lattice).

We briefly summarize the structure of this article. In Section 2 we provide the version of the recursion theorem that we are going to use for the proof of our characterization of effectively discontinuous problems via the discontinuity problem. We also introduce the universal function U and other related concepts. In Section 3 we introduce the discontinuity problem, the notion of effective discontinuity, and we prove related results. In Section 4 we characterize effective discontinuity using Wadge games and we study the relation to determinacy of other games such as Lipschitz games and Gale–Stewart games. In Section 5 we briefly discuss computable discontinuity of characteristic functions and we indicate how this is

³Andretta also proved that Wadge determinacy implies the axiom of countable choice for Baire space [1, Theorem 3], whereas Wadge proved that Wadge determinacy implies the perfect subset property for Baire space [27, Theorem II.C.2].

related to (suitable generalizations of) the notion of productivity. In the conclusions (Section 6) we mention a number of open problems and suggestions for further directions of research.

§2. The universal function and the recursion theorem. We recall that a function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is *computable*, if there is some computable monotone word function $f : \mathbb{N}^* \rightarrow \mathbb{N}^*$ that *approximates* F in the sense that $F(p) = \sup_{w \sqsubseteq p} f(w)$ holds for all $p \in \text{dom}(F)$. Likewise, F is continuous if and only if an analogous condition holds for an arbitrary monotone word function f . Hence, we can define a representation Φ of the set of all continuous functions $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ (with natural domains⁴) by encoding graphs of monotone word functions f into names of F . In other words, if $w : \mathbb{N} \rightarrow \mathbb{N}^*$ is a standard bijective numbering of \mathbb{N}^* , then $p = \langle n_0, k_0 \rangle \langle n_1, k_1 \rangle \dots$ is a name of an extension Φ_p of F if F is approximated by some monotone $f : \mathbb{N}^* \rightarrow \mathbb{N}^*$ with $f(w_{n_i}) = w_{k_i}$ and for each $p \in \text{dom}(F)$ and $n \in \mathbb{N}$ there is some $i \in \mathbb{N}$ with $w_{n_i} \sqsubseteq p$ and $|w_{n_i}| > n$. Here $\langle n, k \rangle := \frac{1}{2}(n+k)(n+k+1) + k$ denotes the usual *Cantor pairing function* for $n, k \in \mathbb{N}$. Intuitively, $F = \Phi_p$ means that p is a listing of a sufficiently large portion of the graph of a monotone function $f : \mathbb{N}^* \rightarrow \mathbb{N}^*$ that approximates F . In order to guarantee that Φ is a total representation, one still needs to clarify how to deal with inconsistent names p , i.e., names for which there is no suitable word function f . Inconsistency can be recognized (i.e., inconsistent names form an open set) and hence one can just consider those p as names of the nowhere defined function. See [29, Definition 3.2.9] for the technical details of such a construction of Φ .

Now we can define a computable *universal function*

$$U : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}, \langle q, p \rangle \mapsto \Phi_q(p),$$

for all $p, q \in \mathbb{N}^{\mathbb{N}}$ [29, Theorem 3.2.16(1)]. Weihrauch [28, Theorems 2.10 and 3.5 and Corollary 2.11] (see also [29, Theorem 3.2.16]) proved the following version of the smn-theorem for the representation Φ that comes in a version for computable and a version for continuous functions.

THEOREM 2.1 (smn-Theorem). *For every computable (continuous) partial function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ there exists a computable (continuous) total function $S : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\Phi_{S(q)}(p) = F\langle q, p \rangle$ for all $p, q \in \mathbb{N}^{\mathbb{N}}$.*

Among other things this result implies that Φ is precomplete. We recall that in general a representation $\delta : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$ of a set X is called *precomplete*, if for every computable $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ there exists a total computable $G : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\delta F(p) = \delta G(p)$ for all $p \in \text{dom}(F)$. In other words, precomplete representations are exactly those under which partial computable functions can be extended to total ones.

Using the smn-theorem one can prove the following uniform version of the recursion theorem along the same lines as the classical recursion theorem. It is an immediate corollary of a more general result due to Kreitz and Weihrauch [13,

⁴We note that for mere cardinality reasons there is no representation of *all* partial continuous $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$, but our representation Φ represents sufficiently many such functions in the sense that it contains an extension of any continuous partial function on Baire space.

Theorem 3.4] (see also [29, Theorem 3.3.20]), which characterizes precomplete representations following Ershov’s characterization of precomplete numberings.

THEOREM 2.2 (Uniform recursion theorem). *There exists a total computable function $T : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\Phi_{T(p)} = \Phi_{\Phi_p T(p)}$ for all $p \in \mathbb{N}^{\mathbb{N}}$ such that Φ_p is total.*

As a corollary of this theorem we obtain the following parameterized version of the recursion theorem that also comes in a version for computable and a version for continuous functions.

COROLLARY 2.3 (Parameterized recursion theorem). *For every computable (continuous) function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ there exists a total computable (continuous) function $R : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $UR(q) = F\langle q, R(q) \rangle$ for all $q \in \mathbb{N}^{\mathbb{N}}$.*

PROOF. We prove the version of the statement for continuous functions. For computable functions one just has to replace the word “continuous” by “computable” in all occurrences. Let $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ be continuous. Then by a double application of the smn-theorem (Theorem 2.1) there is a total continuous $S : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\Phi_{\Phi_{S(q)}(r)}(p) = F\langle p, \langle r, q \rangle \rangle$ for all $p, q, r \in \mathbb{N}^{\mathbb{N}}$. Let $T : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ be the computable function from the recursion theorem (Theorem 2.2). Then $R : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $R(q) := \langle TS(q), q \rangle$ for all $q \in \mathbb{N}^{\mathbb{N}}$ is continuous and satisfies

$$UR(q) = \Phi_{TS(q)}(q) = \Phi_{\Phi_{S(q)}TS(q)}(q) = F\langle q, \langle TS(q), q \rangle \rangle = F\langle q, R(q) \rangle$$

for all $q \in \mathbb{N}^{\mathbb{N}}$. ⊖

In the next section we will use this parameterized version of the recursion theorem to prove our characterization of effectively discontinuous functions.

§3. Effectively discontinuous problems. We now introduce a concept of *computable* (and *effective*) *discontinuity*. These are strengthenings of the concept of discontinuity in the sense that the discontinuity is witnessed by a continuous function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. A computably discontinuous problem f is supposed to have no continuous realizer Φ_q and the computable discontinuity function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ computes for every candidate Φ_q a witnessing input $D(q)$ that shows that the candidate Φ_q fails to realize f on that particular input.

DEFINITION 3.1 (Computable discontinuity). Let (X, δ_X) and (Y, δ_Y) be represented spaces. A problem $f : \subseteq X \rightrightarrows Y$ is called *computably discontinuous* (*effectively discontinuous*) if there is a computable (continuous) $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that for all $q \in \mathbb{N}^{\mathbb{N}}$ we obtain

$$D(q) \in \text{dom}(f\delta_X) \text{ and } \delta_Y\Phi_q D(q) \notin f\delta_X D(q).$$

In this case the function D is called a *discontinuity function* of f .

We emphasize that we consider the condition $\delta_Y\Phi_q D(q) \notin f\delta_X D(q)$ also as satisfied when the left-hand side is undefined. If one does not apply this convention, then one has to write out the condition somewhat more detailed as

$$D(q) \in \text{dom}(f\delta_X) \text{ and } (D(q) \in \text{dom}(\delta_Y\Phi_q) \implies \delta_Y\Phi_q D(q) \notin f\delta_X D(q)).$$

Clearly, every computably discontinuous problem f is effectively discontinuous, and every effectively discontinuous problem is discontinuous. The definition of computable (effective) discontinuity is such that a problem $f : \subseteq X \rightrightarrows Y$ has the respective property if and only if its *realizer version* $f^r : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ with $f^r := \delta_Y^{-1} \circ f \circ \delta_X$ has the property. This implies that it suffices to study the concepts of computable and effective discontinuity on Baire space $\mathbb{N}^{\mathbb{N}}$. We will prove that effective and computable discontinuity can be both characterized in terms of the following discontinuity problem.

DEFINITION 3.2 (Discontinuity problem). We define the *discontinuity problem* by $\text{DIS} : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}, p \mapsto \{q \in \mathbb{N}^{\mathbb{N}} : U(p) \neq q\}$.

We note that DIS is total, i.e., for instances $p \notin \text{dom}(U)$ the problem DIS can provide arbitrary $q \in \mathbb{N}^{\mathbb{N}}$ as solutions. It is a direct consequence of the parameterized recursion theorem from Corollary 2.3 that DIS is computably discontinuous.

PROPOSITION 3.3. *DIS is computably discontinuous.*

PROOF. By Corollary 2.3 there is a computable function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $UD(p) = U\langle p, D(p) \rangle = \Phi_p D(p)$ for all $p \in \mathbb{N}^{\mathbb{N}}$. This function D is hence a computable discontinuity function for DIS. ⊖

We generalize this observation with the following result that is illustrated in the diagram in Figure 1.

THEOREM 3.4 (Effective discontinuity). *Let $f : \subseteq X \rightrightarrows Y$ be a problem. Then:*

- (1) $\text{DIS} \leq_W f \iff f$ is computably discontinuous,
- (2) $\text{DIS} \leq_W^* f \iff f$ is effectively discontinuous.

In both cases one can replace W by its strong counterpart sW .

PROOF. Since $f \equiv_{sW} f^r$ and f is computably (effectively) discontinuous if and only if f^r is, it suffices to prove both statements for problems of type $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$.

(1) “ \implies ” Let $\text{DIS} \leq_W f$ hold via computable $H, K : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$, i.e., $H\langle p, FK(p) \rangle \neq U(p)$ and $H\langle p, FK(p) \rangle$ defined for all $p \in \mathbb{N}^{\mathbb{N}}$ and every realizer F of f . In particular, K is total and $K(p) \in \text{dom}(f)$ for every $p \in \mathbb{N}^{\mathbb{N}}$. By the parameterized recursion theorem from Corollary 2.3 there is some computable $R : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that

$$H\langle R(q), U\langle q, KR(q) \rangle \rangle = UR(q),$$

for all $q \in \mathbb{N}^{\mathbb{N}}$. Then KR is a total computable function and $KR(q) \in \text{dom}(f)$ for all $q \in \mathbb{N}^{\mathbb{N}}$. Let $q \in \mathbb{N}^{\mathbb{N}}$ be such that $KR(q) \in \text{dom}(\Phi_q)$ and let us assume that $U\langle q, KR(q) \rangle = \Phi_q KR(q) \in fKR(q)$. Since f is realizable, there is a realizer F of f with $FKR(q) = U\langle q, KR(q) \rangle$ and hence

$$H\langle R(q), U\langle q, KR(q) \rangle \rangle = H\langle R(q), FKR(q) \rangle \neq UR(q)$$

follows by the choice of H, K in contradiction to the choice of R . Hence the assumption was wrong, i.e., $\Phi_q KR(q) \notin fKR(q)$ and $D := KR$ is a total computable discontinuity function for f . “ \impliedby ” Let f be computably discontinuous with a corresponding computable discontinuity function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. By the

smn-theorem (Theorem 2.1) there is a total computable $R : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $U\langle R(p), q \rangle = U(p)$ for all $p, q \in \mathbb{N}^{\mathbb{N}}$. Now for every realizer F of f and $p \in \mathbb{N}^{\mathbb{N}}$ we obtain $DR(p) \in \text{dom}(f)$ and $FDR(p) \in fDR(p)$ and hence

$$U(p) = U\langle R(p), DR(p) \rangle = \Phi_{R(p)}DR(p) \neq FDR(p),$$

since D is a discontinuity function. Hence, DR is a computable function that witnesses $\text{DIS} \leq_{\text{sw}} f$.

(2) The proof is literally the same as above, except that H, K, R , and D are supposed to be continuous instead of computable. ⊖

We note that the proof of “ \Leftarrow ” shows that we could replace the strong versions of Weihrauch reducibility by an even stronger form of reducibility that only uses the inner reduction function K and no outer reduction function H .

Now we can ask the question whether every discontinuous problem is automatically computably discontinuous, i.e., whether the discontinuity problem DIS is the smallest discontinuous problem with respect to the computable version of Weihrauch reducibility. This is clearly not the case, as the following example shows. In fact, we can infer this from the existence of immune sets. We recall that a set $A \subseteq \mathbb{N}$ is called *immune* if it is infinite but does not contain any infinite c.e. subset.

EXAMPLE 3.5. Let $A \subseteq \mathbb{N}$ and let $f_A : \subseteq [0, 1] \rightarrow \mathbb{R}$ be defined by $\text{dom}(f_A) := \{2^{-n} : n \in A\} \cup \{0\}$ and

$$f_A(x) := \begin{cases} 1 & \text{if } x = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then we obtain:

- (1) f_A discontinuous $\iff A$ infinite.
- (2) f_A computably discontinuous $\iff A$ contains an infinite c.e. subset.
- (3) f_A discontinuous and not computably discontinuous $\iff A$ is immune.

PROOF. (1) is obvious and (3) follows from (1) and (2). For the proof of (2), we note that given a computable discontinuity function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ for f_A , we get an infinite c.e. subset of A as follows. Given $n \in \mathbb{N}$ and $A_n = \{0, 1, 2, \dots, n\}$ we can compute a name q of a realizer Φ_q of the continuous function f_{A_n} , and $D(q)$ has to provide a name of an input $2^{-k} \in [0, 1]$ to f_A on which the realizer Φ_q fails. But this means that $k \in A \setminus A_n$. The collection of all those k for $n = 0, 1, 2, \dots$ forms an infinite c.e. subset of A . On the other hand, given an infinite c.e. subset $B \subseteq A$ and a potential realizer Φ_q of f_A , we can evaluate Φ_q successively on partial names of inputs that correspond to the compact interval $[0, 2^{-k}]$ with $k = 0, 1, 2, \dots$. Then either Φ_q eventually produces an output that excludes 0 or that excludes 1 or neither of this ever happens. In the first case, we can find some $m \in B$ with $m > k$ and produce a name of 2^{-m} as output of $D(q)$, since this is a value on which Φ_q fails to realize $f_A(2^{-m}) = 0$. In the second case, we produce a name of 0 as output of $D(q)$, since this is then a value on which Φ_q fails to realize $f_A(0) = 1$. As long as neither the first nor the second case occurs, we produce the open interval $(-2^{-k}, 2^{-k+1})$ as approximation of an output of $D(q)$. If the first and the second case never occurs, then this produces a name of 0 as output of $D(q)$, which is then not in the domain of Φ_q . Altogether, D is a computable discontinuity function for f_A . ⊖

The following result shows that with respect to the computable version of Weihrauch reducibility we can even get an infinite descending chain of discontinuous problems that are not computably discontinuous. If we restrict the discontinuity problem to the Turing cone $[p] := \{q \in \mathbb{N}^{\mathbb{N}} : p \leq_T q\}$, we obtain an effectively discontinuous problem $\text{DIS}_{|[p]}$. With increasing complexity of p these problems get weaker. By $p^{(n)}$ we denote the n -th Turing jump of $p \in \mathbb{N}^{\mathbb{N}}$.

PROPOSITION 3.6. $\text{DIS}_{|[p^{(n+1)}]} \leq_W \text{DIS}_{|[p^{(n)}]}$ and $\text{DIS}_{|[p^{(n)}]}$ is effectively discontinuous for all $p \in \mathbb{N}^{\mathbb{N}}$ and $n \in \mathbb{N}$, but not computably so for $n \geq 1$.

PROOF. It is clear that $\text{DIS}_{|[p^{(n+1)}]} \leq_{\text{sw}} \text{DIS}_{|[p^{(n)}]}$ holds for all $n \in \mathbb{N}$, as the former problem is a restriction of the latter one. We have $\text{DIS}_{|[p^{(n)}]} \not\leq_W \text{DIS}_{|[p^{(n+1)}]}$, as an instance $q \equiv_T p^{(n)}$ of $\text{DIS}_{|[p^{(n)}]}$ cannot be computably mapped to an instance of $\text{DIS}_{|[p^{(n+1)}]}$. Hence, $\text{DIS}_{|[p^{(n)}]}$ is not computably discontinuous for all $p \in \mathbb{N}^{\mathbb{N}}$ and $n \geq 1$ by Theorem 3.4.

On the other hand, every function Φ_q has names of Turing degree above any $p \in \mathbb{N}^{\mathbb{N}}$, and we can even continuously determine such names by adding redundant information in the code q that encodes p (for instance by repeating the n -th entry of q exactly $p(n)$ times). In other words, for every $p \in \mathbb{N}^{\mathbb{N}}$ there is a continuous function $R_p : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\Phi_q = \Phi_{R_p(q)}$ and $p \leq_T R_p(q)$. Hence $\text{DIS} \leq_W^* \text{DIS}_{|[p]}$ holds for every $p \in \mathbb{N}^{\mathbb{N}}$. That is, $\text{DIS}_{|[p]}$ is effectively discontinuous by Theorem 3.4. \dashv

Hence, we even have an infinite descending chain of effectively discontinuous problems below DIS with respect to the computable version of Weihrauch reducibility. In particular, an effectively discontinuous problem does not need to be computably discontinuous. The next question is whether there are discontinuous problems, which are not even effectively discontinuous. The following proposition provides a sufficient condition for such an example.

PROPOSITION 3.7. Let $\iota : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ be injective and $B := \text{range}(\iota)$. We consider the problem $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ with the domain $\text{dom}(f) := B$ and

$$f(p) := \begin{cases} \mathbb{N}^{\mathbb{N}} \setminus \{\Phi_{\iota^{-1}(p)}(p)\} & \text{if } p \in \text{dom}(\Phi_{\iota^{-1}(p)}), \\ \mathbb{N}^{\mathbb{N}} & \text{otherwise,} \end{cases}$$

for all $p \in B$. Then f is discontinuous. Moreover, if f is effectively discontinuous, then there is a continuous embedding $g : 2^{\mathbb{N}} \hookrightarrow B$.

PROOF. A function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is a discontinuity function for f if and only if $\text{range}(D) \subseteq B$ and

$$D(q) \in \text{dom}(\Phi_q) \implies \Phi_q D(q) = \Phi_{\iota^{-1}D(q)} D(q)$$

holds for all $q \in \mathbb{N}^{\mathbb{N}}$. Clearly, $D = \iota$ is a discontinuity function for f , albeit not necessarily a continuous one. Nevertheless, this shows that f is discontinuous.

Let us now assume that $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is a continuous function with the above property. By the smn-theorem (Theorem 2.1) there exists a computable total function $R : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\Phi_{R(q)}(p) = q$ for all $p, q \in \mathbb{N}^{\mathbb{N}}$. Then we obtain

$$q = \Phi_{R(q)} DR(q) = \Phi_{\iota^{-1}DR(q)} DR(q),$$

for all $q \in \mathbb{N}^{\mathbb{N}}$, which is only possible if $DR : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is injective. In particular, $DR|_{2^{\mathbb{N}}} : 2^{\mathbb{N}} \rightarrow B$ is a continuous embedding of Cantor space into B . \dashv

Hence, the existence of an injective map $\iota : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with a range $B = \text{range}(\iota)$ into which Cantor space cannot be continuously embedded is sufficient to guarantee the existence of a problem f that is discontinuous, but not effectively discontinuous.

For this purpose it is sufficient to show that there exists a set $B \subseteq \mathbb{N}^{\mathbb{N}}$ of continuum cardinality, which violates the perfect subset property. A set B satisfies the *perfect subset property* if it is either countable or it contains a non-empty *perfect* subset, which is a subset that is closed and has no isolated points. We recall that every $A \subseteq \mathbb{N}^{\mathbb{N}}$ into which Cantor space can be continuously embedded contains a perfect subset. It is a well-known fact that there are so-called *Bernstein sets* [3], which violate the perfect subset property, at least if we assume the axiom of choice [12, Exercise 8.24].

FACT 3.8 (Bernstein set). *Assuming AC there exists a Bernstein set $B \subseteq \mathbb{N}^{\mathbb{N}}$, which is a set B such that B as well as its complement $\mathbb{N}^{\mathbb{N}} \setminus B$ has non-empty intersection with every uncountable closed set $A \subseteq \mathbb{N}^{\mathbb{N}}$.*

The construction of a Bernstein set $B \subseteq \mathbb{R}$ provided in [12, Exercise 8.24] works equally well for $B \subseteq \mathbb{N}^{\mathbb{N}}$ and is by transfinite recursion. This construction necessarily requires the axiom of choice. Indeed, if we assume BP, i.e., that every subset $B \subseteq \mathbb{N}^{\mathbb{N}}$ has the Baire property, then no Bernstein set can exist [12] (see also Question 4.19). From the point of view of computability theory Bernstein sets play a similar rôle as immune sets. Actually, together with Proposition 3.7 and Fact 3.8 we directly obtain the following conclusion.

COROLLARY 3.9. *If we assume AC, then there exists a problem $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ that is discontinuous, but not effectively discontinuous.*

Similarly as in the case of Proposition 3.6, one could object that the problem f constructed here is not genuinely less discontinuous than DIS, but only “simpler” as instances are artificially made harder. In other words, the problem f considered here has a very complicated domain, which in the case of our proof of Corollary 3.9 is a Bernstein set.

With the next result we dispel this objection by constructing a total problem $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ that is discontinuous, but not effectively so. Again, the construction is based on the axiom of choice, and we directly perform a transfinite recursion. We can arrange this construction even such that f is parallelizable. We recall that $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ is *parallelizable* if and only if $f \equiv_w \langle f \rangle$, where $\langle f \rangle$ is defined by $\langle f \rangle : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$, $\langle p_0, p_1, \dots \rangle \mapsto \langle f(p_0), f(p_1), \dots \rangle$. By $|X|$ we denote the *cardinality* of a set X .

THEOREM 3.10. *Assuming AC, there exists a total parallelizable $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ that is discontinuous, but not effectively discontinuous.*

PROOF. It suffices to construct a total $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ that is discontinuous and such that $\langle f \rangle : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$, $\langle p_0, p_1, \dots \rangle \mapsto \langle f(p_0), f(p_1), \dots \rangle$ is not effectively discontinuous. Then $\langle f \rangle$ has the desired properties, since it is discontinuous and parallelizable. We note that the set $C(\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}})$ of total continuous functions

$g : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ has continuum cardinality. Hence, by the axiom of choice there is a transfinite enumeration $(r_{\xi})_{\xi < 2^{\aleph_0}}$ of $r_{\xi} \in \mathbb{N}^{\mathbb{N}}$ such that $C(\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}) = \{\Phi_{r_{\xi}} : \xi < 2^{\aleph_0}\}$.

A problem $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ is discontinuous and $\langle f \rangle$ is not effectively discontinuous if the following two requirements are satisfied:

- (1) $(\forall \xi < 2^{\aleph_0})(\exists q \in \mathbb{N}^{\mathbb{N}})(\Phi_q \Phi_{r_{\xi}}(q) \in \langle f \rangle(\Phi_{r_{\xi}}(q)) \text{ and } \Phi_q \text{ total}),$
- (2) $(\forall \xi < 2^{\aleph_0})(\exists p \in \mathbb{N}^{\mathbb{N}}) \Phi_{r_{\xi}}(p) \notin f(p).$

The second condition guarantees that f has no continuous realizer $\Phi_{r_{\xi}}$, and the first condition guarantees that no $\Phi_{r_{\xi}}$ is a discontinuity function for $\langle f \rangle$. We build f by transfinite recursion. For this purpose we construct two increasing sequences $(N_{\xi})_{\xi < 2^{\aleph_0}}$ and $(P_{\xi})_{\xi < 2^{\aleph_0}}$ of sets $N_{\xi}, P_{\xi} \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$. The “negative list” N_{ξ} ensures that f is discontinuous and the “positive list” P_{ξ} ensures that $\langle f \rangle$ is not effectively discontinuous. The construction will be such that $|N_{\xi}| = |\xi|$ and $|P_{\xi}| \leq |\xi| \cdot \aleph_0$. During the construction we frequently use the axiom of choice without further mention. The transfinite recursion goes as follows. We start with $N_0 := P_0 := \emptyset$. For each $0 < \xi < 2^{\aleph_0}$ we first choose some $p \in \mathbb{N}^{\mathbb{N}}$ that does not appear as a first component in $M := \bigcup_{\lambda < \xi} (P_{\lambda} \cup N_{\lambda})$ and we define $N_{\xi} := \{(p, \Phi_{r_{\xi}}(p))\} \cup \bigcup_{\lambda < \xi} N_{\lambda}$. Such a choice is possible, as $|M| \leq |\xi| + |\xi| \cdot \aleph_0 = \max(|\xi|, \aleph_0) < 2^{\aleph_0}$ using the usual rules of cardinal arithmetic [11, Corollary 3.7.8]. Secondly, we also choose some $q \in \mathbb{N}^{\mathbb{N}}$ such that Φ_q is total and such that no $\pi_i \Phi_q \Phi_{r_{\xi}}(q)$ with $i \in \mathbb{N}$ appears in any second component of N_{ξ} . Here $\pi_i : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}, \langle p_0, p_1, \dots \rangle \mapsto p_i$ denotes the projection on the i -th component. Such a choice is possible, as $|N_{\xi}| \leq |\xi| < 2^{\aleph_0}$ and Φ_q for $q \in \mathbb{N}^{\mathbb{N}}$ includes all the constant total functions. We define $P_{\xi} := \{(\pi_i \Phi_{r_{\xi}}(q), \pi_i \Phi_q \Phi_{r_{\xi}}(q)) : i \in \mathbb{N}\} \cup \bigcup_{\lambda < \xi} P_{\lambda}$. The construction guarantees that $|N_{\xi}| = |\xi|$ and $|P_{\xi}| \leq |\xi| \cdot \aleph_0$. This ends the transfinite recursion. We now define $P := \bigcup_{\xi < 2^{\aleph_0}} P_{\xi}$ and $N := \bigcup_{\xi < 2^{\aleph_0}} N_{\xi}$. The construction guarantees that P is the graph of a partial problem $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ and N is the graph of a single-valued function $g : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $N \cap P = \emptyset$. Actually, P guarantees that f satisfies condition (1) and N guarantees that f satisfies condition (2). We still need to extend P to the graph of a total problem $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ without affecting the conditions (1) and (2). For this purpose, we choose for every $p \in \mathbb{N}^{\mathbb{N}}$ that does not yet appear in a first component of P , some $s \in \mathbb{N}^{\mathbb{N}}$ such that $(p, s) \notin N$ and we add (p, s) to P . This is possible since N is the graph of a single-valued function g . Now P is the graph of a suitable total problem $f : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$. ⊖

In Corollary 4.11 we will see that without the axiom of choice (AC) we cannot construct discontinuous problems that are not effectively discontinuous. We close with mentioning that the example in Theorem 3.10 cannot be strengthened to a single-valued function.

PROPOSITION 3.11 (Discontinuous functions). *Every single-valued $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is either continuous or effectively discontinuous.*

PROOF. By $LPO : \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$ we denote the characteristic function of $\{000 \dots\}$. It is well-known that every discontinuous $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ satisfies $LPO \leq_w^* f$ [30, Lemma 8.2.6]. It is easy to see that $DIS \leq_{sw} LPO$ holds (given an input $p \in \mathbb{N}^{\mathbb{N}}$ for the universal function U , use LPO to determine whether $U(p) = 000 \dots$ or not). The statement now follows with Theorem 3.4. ⊖

This result can be extended to functions $f : X \rightarrow Y$ on admissibly represented spaces X, Y (which can be proved similarly as [22, Theorem 4.13]).

§4. A game characterization. In this section we want to characterize continuity and effective discontinuity using games. It is common in descriptive set theory and computability theory to use games to characterize reducibilities and other properties [12, 14, 25] (see [16, 26] for historical surveys).

Wadge [27] introduced games on subsets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ to characterize the reducibility that is named after him. Nobrega and Pauly [20, 21] have used a modification of Wadge games for problems $f : \subseteq X \rightrightarrows Y$ in order to characterize lower cones in the Weihrauch lattice. We consider similar generalized⁵ versions of Wadge and Lipschitz games, defined as follows. We recall that by $f^r : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ we denote the realizer version of a problem $f : \subseteq X \rightrightarrows Y$ (see Section 3).

DEFINITION 4.1 (Wadge game of problems). Let $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be a problem. In a *Wadge game* f two players I and II consecutively play words, with Player I starting:

- Player I: $x_0 \ x_1 \ x_2 \ \dots \ =: x,$
- Player II: $y_0 \ y_1 \ y_2 \ \dots \ =: y,$

with $x_i, y_i \in \mathbb{N}^*$. The concatenated sequences $(x, y) \in (\mathbb{N}^{\mathbb{N}} \cup \mathbb{N}^*)^2$ are called a *run* of the game f . We say that Player II *wins* the run (x, y) of f , if $(x, y) \in \text{graph}(f)$ or $x \notin \text{dom}(f)$. Otherwise Player I *wins*. A Wadge game is called a *Lipschitz game* if $x_i, y_i \in \mathbb{N}$. A *Wadge* or *Lipschitz game* of a general problem $f : \subseteq X \rightrightarrows Y$ is understood to be the corresponding game of the realizer version $f^r : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$.

We allow both players to play arbitrary words, including the empty word. One can see that Player I does not take any advantage of playing words and he could be restricted to natural numbers in a Wadge game, without loss of generality. Likewise, Player II does not take any advantage of playing arbitrary words, it would suffice to allow numbers and the empty word, where the empty word essentially corresponds to skipping the corresponding move. This shows that our notion of a Wadge game for problems corresponds to the one of Nobrega and Pauly [20, 21]. For simplicity we have allowed arbitrary words for both players. This does not only lead to a more symmetric definition, but it also simplifies the proof of Theorem 4.3 below. As usual we define winning strategies for games to be word functions that determine moves for one player depending on the moves of the other player.

DEFINITION 4.2 (Winning strategy). Let $f : \subseteq X \rightrightarrows Y$ be a problem and let $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ be a function. We consider the Wadge game f .

- (1) σ is called a *winning strategy* for Player II in the game f , if Player II wins every run of f with her moves being determined by

$$y_i := \sigma(x_0, \dots, x_i),$$

while Player I plays $x_0, x_1, \dots \in \mathbb{N}^*$.

⁵We warn the reader that the extension of the notion of a Wadge game from sets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ to problems does not automatically mean that determinacy properties carry over.

- (2) σ is called a *winning strategy* for Player I in the game f , if Player I wins every run of f with his moves being determined by

$$x_i := \sigma(y_0, \dots, y_{i-1}),$$

while Player II plays $y_0, y_1, \dots \in \mathbb{N}^*$.

Winning strategies for Lipschitz games are defined analogously with functions of type $\sigma : \mathbb{N}^* \rightarrow \mathbb{N}$.

Now our main observation on Wadge games is that winning of Player II characterizes continuity of the problem and winning of Player I effective discontinuity. Nobrega and Pauly have proved a general version of the first observation for lower cones in the Weihrauch lattice [21, Theorem 3.3]. We use some bijective standard numbering $w : \mathbb{N} \rightarrow \mathbb{N}^*$ and we use the notation $\bar{v} := w^{-1}(v)$ for all $v \in \mathbb{N}^*$.

THEOREM 4.3 (Wadge games). *We consider the Wadge game of a given problem $f : \subseteq X \rightrightarrows Y$. Then the following hold:*

- (1) f is continuous \iff Player II has a winning strategy for f ,
- (2) f is effectively discontinuous \iff Player I has a winning strategy for f .

PROOF. Since the Wadge game of f is the Wadge game of f^r and $f^r \equiv_{\text{sw}} f$, it suffices by Theorem 1.4 to consider problems of type $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$.

(1) If f is continuous, then f has a continuous realizer $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$, which is approximated by a monotone function $h : \mathbb{N}^* \rightarrow \mathbb{N}^*$ in the sense that $F(p) = \sup_{w \sqsubseteq p} h(w)$ for all $p \in \text{dom}(F) \supseteq \text{dom}(f)$. Given the moves $w_0, w_1, \dots \in \mathbb{N}^*$ of Player I, we can inductively define the moves $v_i \in \mathbb{N}^*$ by $v_0 \dots v_i := h(w_0 \dots w_i)$ for all $i \in \mathbb{N}$, since h is monotone. Then $\sigma(\varepsilon) := \varepsilon$ and $\sigma(w_0, \dots, w_i) := v_i$ provides a winning strategy $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ for Player II. This is because if $r := w_0 w_1 \dots \in \text{dom}(f)$, then $F(r) = v_0 v_1 \dots \in f(r)$.

Vice versa, let $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ be a winning strategy for Player II. Then we can define $h : \mathbb{N}^* \rightarrow \mathbb{N}^*$ by $h(\varepsilon) := \varepsilon$ and $h(a_0 \dots a_i) := v_0 \dots v_i$ for all $a_0, a_1, \dots \in \mathbb{N}$, where we inductively choose $v_i := \sigma(a_0, \dots, a_i)$. Let $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ be given by $F(p) := \sup_{w \sqsubseteq p} h(w)$. Given an input $r := a_0 a_1 \dots \in \text{dom}(f)$, we obtain $F(r) = v_0 v_1 \dots$ such that $(r, F(r)) \in \text{graph}(f)$, since σ is a winning strategy for Player II. Hence F is a continuous realizer for f .

(2) If f is effectively discontinuous, then there is a continuous $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ that witnesses the discontinuity of f in the sense that $D(p) \in \text{dom}(f)$ and $\Phi_p D(p) \notin f D(p)$ for all $p \in \mathbb{N}^{\mathbb{N}}$. Let $h : \mathbb{N}^* \rightarrow \mathbb{N}^*$ be a monotone function that approximates D , in the sense that $D(p) = \sup_{w \sqsubseteq p} h(w)$ for all $p \in \mathbb{N}^{\mathbb{N}}$. Given the moves $v_0, v_1, \dots \in \mathbb{N}^*$ of Player II, we can inductively define the moves w_i by $w_0 := h(\varepsilon)$ and $w_0 \dots w_i := h(\langle \bar{w}_0, \bar{v}_0 \rangle \dots \langle \bar{w}_0 \dots w_{i-1}, \bar{v}_0 \dots v_{i-1} \rangle)$. We note that $r := w_0 w_1 \dots$ is an infinite sequence $r \in \text{dom}(f)$, since D is total and $\text{range}(D) \subseteq \text{dom}(f)$. And $p := \langle \bar{w}_0, \bar{v}_0 \rangle \langle \bar{w}_0 w_1, \bar{v}_0 v_1 \rangle \dots \in \mathbb{N}^{\mathbb{N}}$ is the name of a function Φ_p with $D(p) = r$. If $q := v_0 v_1 \dots$ is finite, then clearly $(r, q) \notin \text{graph}(f)$. Otherwise, $q = \Phi_p(r) = \Phi_p D(p) \notin f D(p) = f(r)$ and hence also $(r, q) \notin \text{graph}(f)$. This means that $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ with $\sigma(v_0, \dots, v_{i-1}) := w_i$ is a winning strategy for Player I.

Vice versa let $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ be a winning strategy for Player I. We need to define a continuous discontinuity function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ for f . Given $p \in \mathbb{N}^{\mathbb{N}}$, we can determine a monotone $h : \mathbb{N}^* \rightarrow \mathbb{N}^*$ that approximates Φ_p , i.e., such that $\Phi_p(q) = \sup_{w \sqsubseteq q} h(w)$ for all $q \in \text{dom}(\Phi_p)$. We let $D(p) := w_0 w_1 \dots$ where the w_i are inductively given by $w_i := \sigma(v_0, \dots, v_{i-1})$ and $v_0 \dots v_i := h(w_0 \dots w_i)$. Since σ is a winning strategy for Player I, we have $D(p) \in \text{dom}(f)$. In particular, D is total and continuous. Moreover, with $q := v_0 v_1 \dots$ we have $(D(p), q) \notin \text{graph}(f)$. This could mean that q is finite and hence $D(p) \notin \text{dom}(\Phi_p)$ or otherwise $\Phi_p D(p) = q$. In any case, $\Phi_p D(p) \notin f D(p)$ holds and D is a discontinuity function for f . \dashv

In passing, we note that the proof is fully constructive in the sense that computable winning strategies translate into computable functions and vice versa, as stated in the following.

COROLLARY 4.4 (Wadge games). *We consider the Wadge game of a given problem $f : \subseteq X \rightrightarrows Y$. Then the following hold:*

- (1) f is computable \iff Player II has a computable winning strategy for f ,
- (2) f is computably discontinuous \iff Player I has a computable winning strategy for f .

One reason why it is useful to have characterizations of continuity and effective discontinuity in game form is that for certain games determinacy conditions are known and well understood. The axiom of determinacy (AD), which was introduced by Mycielski and Steinhaus [15, 17] states that every Gale–Stewart game is determined. We note that this axiom is inconsistent with the axiom of choice. We recall the definition of Gale–Stewart games [25].⁶

DEFINITION 4.5 (Gale–Stewart game). Let $A \subseteq \mathbb{N}^{\mathbb{N}}$. Then in a Gale–Stewart game A two players I and II consecutively play numbers

- Player I: $x_0 \ x_1 \ x_2 \ \dots =: x$,
- Player II: $y_0 \ y_1 \ y_2 \ \dots =: y$,

with $x_i, y_i \in \mathbb{N}$. The concatenated sequence $r = \langle x, y \rangle \in \mathbb{N}^{\mathbb{N}}$ is called a *run* of the game A . We say that Player II *wins* the run r of A , if $r \in A$. Otherwise Player I *wins*.

What we have described as the Gale–Stewart game of A is usually considered as the Gale–Stewart game of the complement $\mathbb{N}^{\mathbb{N}} \setminus A$. We prefer the complementary version as it fits better to our definition of Wadge games.

Winning strategies for Gale–Stewart games can be defined analogously to Lipschitz games. In fact, Lipschitz games of problems f are essentially Gale–Stewart games on $\text{graph}(f)$, at least for total problems on Baire space. For general problems, the difference can be expressed using totalizations of problems, a concept that was studied in [8, 19].

DEFINITION 4.6 (Totalization). Let $f : \subseteq X \rightrightarrows Y$ be a problem. Then the *totalization* $\top f : X \rightrightarrows Y$ is defined by

$$\top f(x) := \begin{cases} f(x) & \text{if } x \in \text{dom}(f), \\ Y & \text{otherwise.} \end{cases}$$

⁶Such games over binary digits were already considered by Ulam [26].

For a subset $A \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ we use the notation $\langle A \rangle := \{ \langle p, q \rangle : (p, q) \in A \}$. Now we can express the relation between Lipschitz games and Gale–Stewart games on graphs as follows.

PROPOSITION 4.7 (Lipschitz games and Gale–Stewart games). *Let $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be a problem. Then the winning strategies for either player in the Lipschitz game f are identical to the winning strategies of the corresponding player for the Gale–Stewart game $\langle \text{graph}(\top f) \rangle$.*

PROOF. The proof follows from the easy observation that

$$\text{graph}(\top f) = \text{graph}(f) \cup ((\mathbb{N}^{\mathbb{N}} \setminus \text{dom}(f)) \times \mathbb{N}^{\mathbb{N}}),$$

and this is exactly the payoff set for Player II in the Lipschitz game f . ⊣

We note that $f \leq_w \top f$, but in general $\top f$ is not Weihrauch equivalent to f [8].

For general Wadge games the situation is somewhat more subtle as players can play empty words. We can, however, bridge the step between Wadge and Lipschitz games by coding words in numbers. We consider a canonical bijective standard numbering $w : \mathbb{N} \rightarrow \mathbb{N}^*$, $i \mapsto w_i$ of words over natural numbers, which we lift to a partial function

$$w : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}, p \mapsto w_{p(0)} w_{p(1)} w_{p(2)} \dots$$

that we also denote by w . Here $\text{dom}(w) = \{ p \in \mathbb{N}^{\mathbb{N}} : w_{p(0)} w_{p(1)} w_{p(2)} \dots \text{ is infinite} \}$ and hence w is not total (since there is a number n that encodes the empty word). Now we can consider the problem $f^w := w^{-1} \circ f \circ w$, which is f lifted to numbers (that encode words). We note that f^w is not total, even if f is total, since w is not total. We mention that $w : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ can be seen as a precomplete representation of $\mathbb{N}^{\mathbb{N}}$ (in fact, as an alternative way to define the *precompletion* of $\text{id} : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$, as studied in [7, 8]). Since w is computably equivalent to id as a representation, we obtain $f \equiv_{\text{sw}} f^w$. Now we can express our observation as follows.

PROPOSITION 4.8 (Wadge games and Lipschitz games). *Let $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be a problem and let $P \in \{\text{I}, \text{II}\}$. Then the following are equivalent.*

- (1) *Player P has a winning strategy for the Wadge game f .*
- (2) *Player P has a winning strategy for the Lipschitz game f^w .*

PROOF. Given a winning strategy $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ for Player P of the Wadge game f , we have to convert this strategy into a strategy $\lambda : \mathbb{N}^* \rightarrow \mathbb{N}$ for Player P of the Lipschitz game f^w . For this purpose we just have to define λ such that $\lambda(n_0, \dots, n_k) := w^{-1} \sigma(w_{n_0}, \dots, w_{n_k})$. Vice versa, if $\lambda : \mathbb{N}^* \rightarrow \mathbb{N}$ is a winning strategy for Player P of the Lipschitz game f^w , then $\sigma(w_0, \dots, w_k) := w_{\lambda(w^{-1}(w_0), \dots, w^{-1}(w_k))}$ defines a winning strategy $\sigma : \mathbb{N}^{**} \rightarrow \mathbb{N}^*$ for Player P of the Wadge game f . ⊣

If we combine Propositions 4.7 and 4.8 then we obtain the following result.

COROLLARY 4.9 (Wadge games and Gale–Stewart games). *Let $f : \subseteq X \rightrightarrows Y$ be a problem and let $P \in \{\text{I}, \text{II}\}$. Then the following are equivalent.*

- (1) *Player P has a winning strategy for the Wadge game f .*
- (2) *Player P has a winning strategy for the Lipschitz game f^{rw} .*
- (3) *Player P has a winning strategy for the Gale–Stewart game $\langle \text{graph}(\top(f^{rw})) \rangle$.*

We note that $T(f^{rw}) \equiv_{sw} \bar{f}$, where \bar{f} denotes the so-called *completion* of f (see [8, Lemma 4.9]). Hence, the operation of precompletion moves us from Wadge games to Lipschitz games and the operation of completion to Gale–Stewart games.

In order to have some simple terminology at hand, we introduce the following notions of determinacy.

DEFINITION 4.10 (Determinacy). We call a problem $f : \subseteq X \rightrightarrows Y$ *determined*, if it is either continuous or effectively discontinuous. By *Weihrauch determinacy* we understand the property that every problem f is determined.

Hence, by Theorem 3.4 f is determined if either $f \leq_w^* \text{id}$ or $\text{DIS} \leq_w^* f$ holds and by Theorem 4.3 this is the case if and only if the Wadge game f is determined in the sense that either Player I or Player II has a winning strategy. Weihrauch determinacy means that every f satisfies the above dichotomies.

Since the axiom of determinacy AD states that every Gale–Stewart game is determined, i.e., either Player I or Player II has a winning strategy, we immediately get the following conclusion of Corollary 4.9 with the help of Theorem 3.4 (see also [21, Corollary 3.7]).

COROLLARY 4.11 (Determinacy). $ZF + DC + AD$ implies *Weihrauch determinacy*.

We can conclude more from Corollary 4.9. Namely, in $ZF + AC$ it is known by a Theorem of Martin [12, Theorem 20.5] that every Borel set $A \subseteq \mathbb{N}^{\mathbb{N}}$ is determined, i.e., either Player I or Player II has a winning strategy (see also [21, Corollary 3.6]).

PROPOSITION 4.12 (Borel determinacy). *In ZFC every problem $f : \subseteq X \rightrightarrows Y$, such that $\text{graph}(f^r) \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ and $\text{dom}(f^r) \subseteq \mathbb{N}^{\mathbb{N}}$ are Borel sets, is determined.*

PROOF. Firstly, for a problem $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ we note that

$$\text{graph}(f^w) = \text{graph}(w^{-1} \circ f \circ w) = (w \times w)^{-1}(\text{graph}(f))$$

and $\text{dom}(f^w) = \text{dom}(f \circ w) = w^{-1}(\text{dom}(f))$. Since w is continuous with a Π_2^0 -domain $\text{dom}(w)$, it follows that $\text{graph}(f^w)$ and $\text{dom}(f^w)$ are Borel measurable, if $\text{graph}(f)$ and $\text{dom}(f)$ are so. Hence, $\langle \text{graph}(Tf^w) \rangle$ is Borel measurable in this situation. For a general problem $f : \subseteq X \rightrightarrows Y$, we can apply the previous considerations to $f^r : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ and we obtain the claim using Borel determinacy and Corollary 4.9. ⊣

That is even under the axiom of choice AC examples of problems that are discontinuous but not effectively so have to be rather complicated. As Polish spaces admit continuous and total versions of the Cauchy representation (see, e.g., [4, Corollary 4.4.12]), we also obtain the following version of the previous corollary.

COROLLARY 4.13 (Borel determinacy on Polish spaces). *Let X, Y be Polish spaces. In ZFC every problem $f : \subseteq X \rightrightarrows Y$, such that $\text{graph}(f) \subseteq X \times Y$ and $\text{dom}(f) \subseteq X$ are Borel sets, is determined.*

PROOF. For Polish spaces X and Y we can assume that the Cauchy representations $\delta_X : \mathbb{N}^{\mathbb{N}} \rightarrow X$ and $\delta_Y : \mathbb{N}^{\mathbb{N}} \rightarrow Y$ are total [4, Corollary 4.4.12]. They are also continuous maps. Since $f^r = \delta_Y^{-1} \circ f \circ \delta_X$, this implies that $\text{graph}(f^r) = (\delta_X \times \delta_Y)^{-1}(\text{graph}(f))$ and $\text{dom}(f^r) = \delta_X^{-1}(\text{dom}(f))$ are Borel sets, if $\text{graph}(f)$ and $\text{dom}(f)$ are Borel sets. Now the claim follows with Proposition 4.12. ⊣

It is an interesting questions whether we get the inverse implication in Corollary 4.11 in the following sense.

QUESTION 4.14. *Does AD follow from Weihrauch determinacy in ZF + DC?*

We note that a similar question, namely whether Wadge determinacy (in the usual sense of Wadge games for sets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$) implies AD is non-trivial and the answer is not known [1, 2]. For our generalized Wadge games the situation seems to be simpler and we can easily obtain AD from Lipschitz determinacy of problems (with a non-constructive proof).

PROPOSITION 4.15. *In ZF + DC the following are equivalent:*

- (1) *The axiom of determinacy AD.*
- (2) *Every Lipschitz game $f : \subseteq X \rightrightarrows Y$ is determined, i.e., either Player I or Player II has a winning strategy.*

PROOF. That (1) implies (2) was proved in Proposition 4.7. We still need to prove that (2) implies (1). For every set $G \subseteq \mathbb{N}^{\mathbb{N}}$ we define the problem $f_G : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ with $f_G(p) := \{q \in \mathbb{N}^{\mathbb{N}} : \langle p, q \rangle \in G\}$ and $\text{dom}(f_G) := \{p \in \mathbb{N}^{\mathbb{N}} : (\exists q \in \mathbb{N}^{\mathbb{N}}) \langle p, q \rangle \in G\}$. We call G total, if $\text{dom}(f_G) = \mathbb{N}^{\mathbb{N}}$. It is clear that for non-total G Player I in the Gale–Stewart game G always has a winning strategy, and he just needs to play some $p \notin \text{dom}(f_G)$. For total G we obtain $\langle \text{graph}(\top f_G) \rangle = \langle \text{graph}(f_G) \rangle = G$. Hence, by Proposition 4.7 G is determined if the Lipschitz game f_G is so. This proves determinacy of G for every $G \subseteq \mathbb{N}^{\mathbb{N}}$. \dashv

In order to answer Question 4.14 positively one still needs to bridge the gap between Lipschitz determinacy and Wadge determinacy for problems f (possibly using the techniques from [1, 2] or a simpler argument).

We prove that Weihrauch determinacy implies at least Wadge determinacy, in the original sense of Wadge games for sets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$. We can simulate Wadge games for sets A, B using a particular problem $\frac{B}{A}$ (which was introduced in [21]).

DEFINITION 4.16 (Wadge game for sets). Let $A, B \subseteq \mathbb{N}^{\mathbb{N}}$. Then the Wadge game (A, B) is the Wadge game of the problem $\frac{B}{A} : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ with

$$\text{graph} \left(\frac{B}{A} \right) := (A \times B) \cup ((\mathbb{N}^{\mathbb{N}} \setminus A) \times (\mathbb{N}^{\mathbb{N}} \setminus B)).$$

In the Wadge game $\frac{B}{A}$ Player II wins if the corresponding run (x, y) satisfies $x \in A \iff y \in B$ and otherwise Player I wins. This shows that the Wadge game (A, B) is the usual one and as usually, by *Wadge determinacy* we understand the property that every Wadge game (A, B) is determined, i.e., either Player I or Player II has a winning strategy. This immediately yields the following.

COROLLARY 4.17 (Wadge determinacy). *In ZF + DC Weihrauch determinacy implies Wadge determinacy.*

Another conclusion that we can either directly draw from Proposition 3.7 without further ado or from Corollary 4.17 with the help of [27, Theorem II.C.2] is the following.

COROLLARY 4.18 (Perfect subset property). *In ZF + DC Weihrauch determinacy implies that every set $A \subseteq \mathbb{N}^{\mathbb{N}}$ satisfies the perfect subset property.*

If Question 4.14 has a negative answer or cannot be answered easily, then one could ask other questions such as the following.

QUESTION 4.19. *Does the Baire property BP follow from Weihrauch determinacy in ZF + DC?*

By the Baire property BP we mean that statement that every subset $A \subseteq \mathbb{N}^{\mathbb{N}}$ can be written as symmetric difference $A = U \Delta M$ with an open set $U \subseteq \mathbb{N}^{\mathbb{N}}$ and a meager set $M \subseteq \mathbb{N}^{\mathbb{N}}$. If the answer to this question is also negative, then one can ask the following modified version of Question 4.14.

QUESTION 4.20. *Does AD follow from Weihrauch determinacy in ZF + DC + BP?*

A positive answer to this question is not completely unlikely in light of Proposition 4.15, given that Wadge determinacy and Lipschitz determinacy (both in the usual sense of games for sets $A, B \subseteq \mathbb{N}^{\mathbb{N}}$) have been proven to be equivalent in ZF + DC + BP by Andretta [1, 2].

§5. Computable discontinuity and productivity. In this section we briefly want to discuss the question what computable discontinuity means for subsets and how the notion is linked to the notion of productivity. Here, for every subset $A \subseteq X$ of some fixed space X , we denote by

$$\chi_A : X \rightarrow \mathbb{S}, x \mapsto \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{otherwise} \end{cases}$$

the *characteristic function* of A . The codomain $\mathbb{S} = \{0, 1\}$ is Sierpiński space that is equipped with the total representation $\delta_{\mathbb{S}}$ with $\delta_{\mathbb{S}}(p) = 0 : \iff p = 000\dots$. It is well-known that for any represented space X and $A \subseteq X$ the characteristic function χ_A is continuous if and only if A is open.

As a side remark we mention that the characteristic function in this form can also nicely be used to embed Wadge reducibility and many-one reducibility into the strong version of Weihrauch reducibility for non-trivial sets. We recall that for $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ the set A is called *Wadge reducible* to B , in symbols $A \leq_{\text{sw}} B$, if there is a continuous $f : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $A = f^{-1}(B)$. Likewise, we say for $A, B \subseteq \mathbb{N}$ that A is *many-one reducible* to B , in symbols $A \leq_{\text{m}} B$, if there is a computable $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $A = f^{-1}(B)$. Since the only continuous functions $f : \mathbb{S} \rightarrow \mathbb{S}$ are the identity and the two constant functions, we directly get the following characterization of Wadge and many-one reducibility in terms of the topological version of strong Weihrauch reducibility.

PROPOSITION 5.1 (Reducibilities). *We obtain*

- (1) $\chi_A \leq_{\text{sw}} \chi_B \iff A \leq_{\text{m}} B$ for $A, B \subseteq \mathbb{N}$ with $B \notin \{\emptyset, \mathbb{N}\}$.
- (2) $\chi_A \leq_{\text{sw}}^* \chi_B \iff A \leq_{\text{w}} B \iff \frac{B}{A} \leq_{\text{w}}^* \text{id}$ for $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ with $B \notin \{\emptyset, \mathbb{N}^{\mathbb{N}}\}$.

Hence, the corresponding reducibility structures can be embedded into the corresponding strong versions of the Weihrauch lattice for non-trivial sets. By $\mathcal{O}(X)$ we denote the set of open subsets of X equipped with the representation $\delta_{\mathcal{O}(X)}$, defined by $\delta_{\mathcal{O}(X)}(p) = U : \iff \Phi_p$ is a realizer of χ_U . Now one can ask what it means for χ_A to be computably discontinuous. The following result answers this question for Baire space $X = \mathbb{N}^{\mathbb{N}}$ using the *symmetric difference*

problem $\Delta_A : \subseteq \mathcal{O}(X) \Rightarrow X, U \mapsto A\Delta U$. Here $A\Delta U := (A \setminus U) \cup (U \setminus A)$ denotes the symmetric difference of $A, U \subseteq X$. The problem Δ_A is total for non-open $A \subseteq X$.

PROPOSITION 5.2 (Symmetric difference). *Let $A \subseteq \mathbb{N}^{\mathbb{N}}$ not be open. Then the problem Δ_A is computable if and only if χ_A is computably discontinuous.*

PROOF. If $\chi_A : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{S}$ is computably discontinuous, then there is a computable discontinuity function $D : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ for χ_A . Given a name q of a realizer Φ_q of the continuous function $\chi_U : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{S}$ for any given $U \in \mathcal{O}(\mathbb{N}^{\mathbb{N}})$, the value $D(q)$ is a name for some point $x \in X$ with $\chi_U(x) \neq \chi_A(x)$, i.e., $x \in A\Delta U$. That is, Δ_A is realized by D and hence computable. On the other hand, let Δ_A be computable and $A \subseteq \mathbb{N}^{\mathbb{N}}$ not open. Given a name q of some potential realizer Φ_q of χ_A , we can convert q computably into a name $r = G(q)$ of some total function Φ_r with $\delta_{\mathbb{S}}\Phi_r = \delta_{\mathbb{S}}\Phi_q$, because $\delta_{\mathbb{S}}$ is precomplete (the idea is that Φ_r produces zero output as long as Φ_q makes no other information available). Let us denote by $G : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ the corresponding computable function. Now there is some open $U \subseteq \mathbb{N}^{\mathbb{N}}$ such that Φ_r is a realizer of χ_U , namely $U := \Phi_r^{-1}(\mathbb{N}^{\mathbb{N}} \setminus \{000\dots\}) = \Phi_r^{-1}\delta_{\mathbb{S}}^{-1}(\{1\})$. Since A is not open, $\Delta_A(U)$ is defined and non-empty. Let F be a computable realizer of Δ_A and $D := FG$. Then $F(r) \in \Delta_A(U) = (A \setminus U) \cup (U \setminus A)$, i.e., $\chi_U D(q) \neq \chi_A D(q)$. This implies $\delta_{\mathbb{S}}\Phi_q D(q) \neq \chi_A D(q)$, since either $D(q) \notin \text{dom}(\Phi_q)$ or otherwise $\delta_{\mathbb{S}}\Phi_q D(q) = \delta_{\mathbb{S}}\Phi_r D(q) = \chi_U D(q)$. That is D is a computable discontinuity function of χ_A . ⊣

An analogous statement holds if we replace computable by continuous in both occurrences. However, in this case the statement is void as any χ_A for non-open A is effectively discontinuous by (a suitable extension of) Proposition 3.11 or by observing that Δ_A for non-open A is always continuous. We can also summarize Proposition 5.2 as follows.

COROLLARY 5.3 (Symmetric difference). $\Delta_A \leq_w \text{id} \iff \text{DIS} \leq_w \chi_A$ for non-open $A \subseteq \mathbb{N}^{\mathbb{N}}$.

Proposition 5.2 shows that the notion of computable discontinuity is also formally related to the notion of productivity. Weihrauch has introduced a topological and a computability-theoretic notion of productivity for subsets $A \subseteq \mathbb{N}^{\mathbb{N}}$ [28, Definition 4.4], [29, Definition 3.2.24]. If, in a similar way, we transfer the definition of completely productive sets, as originally defined by Dekker [10], then we could say that non-open $A \subseteq \mathbb{N}^{\mathbb{N}}$ is completely productive if Δ_A is computable. Using this terminology, non-open $A \subseteq \mathbb{N}^{\mathbb{N}}$ is completely productive if and only if χ_A is computably discontinuous by Proposition 5.2. Myhill [18] proved that a set $A \subseteq \mathbb{N}$ is productive if and only if A is completely productive (see also the proofs in [24, Theorem VII, Section 11.3], [29, Corollary 2.6.8]). It is clear that complete productivity for $A \subseteq \mathbb{N}^{\mathbb{N}}$ implies productivity. We leave it as a task to the reader to study whether productivity also implies complete productivity for Baire space $\mathbb{N}^{\mathbb{N}}$ (or even more general spaces).

§6. Conclusions. We have introduced the discontinuity problem DIS and we have provided some evidence that one can consider it as the simplest natural unsolvable problem with respect to the topological version of Weihrauch reducibility. At least

in $ZF + DC + AD$ it turns out that it actually induces the minimal discontinuous Weihrauch degree.

More results on the discontinuity problem will be provided in a forthcoming article [5]. While the original definition of DIS is in terms of a universal function U , it is useful to have a characterization in purely set-theoretic terms. In [5] we prove that the discontinuity problem is equivalent to the *range non-equality problem* defined by $NRNG : \mathbb{N}^{\mathbb{N}} \rightrightarrows 2^{\mathbb{N}}, p \mapsto \{A \in 2^{\mathbb{N}} : A \neq \text{range}(p - 1)\}$. Here for $p \in \mathbb{N}^{\mathbb{N}}$ the finite or infinite sequence $p - 1 \in \mathbb{N}^{\mathbb{N}} \cup \mathbb{N}^*$ is the sequence that is obtained as concatenation of $p(0) - 1, p(1) - 1, p(2) - 1, \dots$, where $- 1$ is identified with the empty word ε .

PROPOSITION 6.1. $DIS \equiv_{sW} NRNG$.

In [5] we also discuss algebraic properties of the discontinuity problem DIS and one important property is that the parallelization \widehat{DIS} of the discontinuity problem is equivalent to the non-computability problem NON.

THEOREM 6.2 (Non-computability is parallelized discontinuity). $NON \equiv_{sW} \widehat{DIS}$.

The *non-computability problem* is defined with the help of Turing reducibility \leq_T by $NON : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}, p \mapsto \{q \in \mathbb{N}^{\mathbb{N}} : q \not\leq_T p\}$. This result supports the slogan that “non-computability is the parallelization of discontinuity” and underlines that the discontinuity problem is a natural one. The discontinuity problem can itself be obtained by stashing (a dual operation to parallelization introduced in [5]) starting from other natural problems such as LPO, LLPO, etc. Hence, it is nicely related in an algebraic way to other natural problems in the Weihrauch lattice.

Acknowledgments. We would like to thank Matthias Schröder for a discussion on the relevance of the axiom of determinacy and different forms of the axiom of choice for computable analysis. Likewise, we would like to thank Arno Pauly for a discussion on Wadge games for problems that has helped to improve the corresponding results. This work has been supported by the *National Research Foundation of South Africa* (Grant Number 115269).

REFERENCES

- [1] A. ANDRETTA, *Equivalence between Wadge and Lipschitz determinacy*. *Annals of Pure and Applied Logic*, vol. 123 (2003), nos. 1–3, pp. 163–192.
- [2] ———, *More on Wadge determinacy*. *Annals of Pure and Applied Logic*, vol. 144 (2006), nos. 1–3, pp. 2–32.
- [3] F. BERNSTEIN, *Zur Theorie der trigonometrischen Reihe*. *Sitzungsberichte der Königlich Sächsischen Gesellschaft der Wissenschaften zu Leipzig, Math.-Phys. Klasse*, vol. 60 (1908), pp. 325–338.
- [4] V. BRATTKA, *Recursive and computable operations over topological structures*, Ph.D. thesis, Department of Computer Science, University of Hagen, 1998.
- [5] ———, *Stashing-parallelization pentagons*. *Logical Methods in Computer Science*, vol. 17 (2021), no. 4, pp. 20:1–20:29.
- [6] V. BRATTKA, D. D. DZHAFAROV, A. MARCONE, and A. PAULY, editors, *Measuring the complexity of computational content: From combinatorial problems to analysis*. *Dagstuhl Reports*, vol. 8, Dagstuhl, Germany, Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2019.
- [7] V. BRATTKA and G. GHERARDI, *Weihrauch goes Brouwerian*, this JOURNAL, vol. 85 (2020), no. 4, pp. 1614–1653.
- [8] ———, *Completion of choice*. *Annals of Pure and Applied Logic*, vol. 172 (2021), no. 3, p. 102914.

- [9] V. BRATTKA, G. GHERARDI, and A. PAULY, *Weihrauch complexity in computable analysis*, **Handbook of Computability and Complexity in Analysis** (V. Brattka and P. Hertling, editors), Theory and Applications of Computability, Springer, Cham, 2021, pp. 367–417.
- [10] J. C. E. DEKKER, *Productive sets*. **Transactions of the American Mathematical Society**, vol. 78 (1955), pp. 129–149.
- [11] K. DEVLIN, *The Joy of Sets, Fundamentals of Contemporary Set Theory*, second ed., Undergraduate Texts in Mathematics, Springer, New York, 1993.
- [12] A. S. KECHRIS, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, vol. 156, Springer, Berlin, 1995.
- [13] C. KREITZ and K. WEIHRAUCH, *Theory of representations*. **Theoretical Computer Science**, vol. 38 (1985), pp. 35–53.
- [14] Y. N. MOSCHOVAKIS, *Descriptive Set Theory*, second ed., Mathematical Surveys and Monographs, vol. 155, American Mathematical Society, Providence, RI, 2009.
- [15] J. MYCIELSKI, *On the axiom of determinateness*. **Fundamenta Mathematicae**, vol. 53 (1963/64), pp. 205–224.
- [16] ———, *Games with perfect information*, **Handbook of Game Theory with Economic Applications**, vol. I (R. J. Aumann and S. Hart, editors), Handbooks in Economics, vol. 11, North-Holland, Amsterdam, 1992, pp. 41–70.
- [17] J. MYCIELSKI and H. STEINHAUS, *A mathematical axiom contradicting the axiom of choice*. **Bulletin L'Académie Polonaise des Sciences, Série des Sciences Mathématiques, Astronomiques et Physiques**, vol. 10 (1962), pp. 1–3.
- [18] J. MYHILL, *Creative sets*. **Zeitschrift für Mathematische Logik und Grundlagen der Mathematik**, vol. 1 (1955), pp. 97–108.
- [19] E. NEUMANN and A. PAULY, *A topological view on algebraic computation models*. **Journal of Complexity**, vol. 44 (2018), Suppl. C, pp. 1–22.
- [20] H. NOBREGA, *Games for functions—Baire classes, Weihrauch degrees, transfinite Computations, and ranks*. Ph.D. thesis, Institute for Logic, Language and Computation, Universiteit van Amsterdam, 2018.
- [21] H. NOBREGA and A. PAULY, *Game characterizations and lower cones in the Weihrauch degrees*. **Logical Methods in Computer Science**, vol. 15 (2019), no. 3, Article no. 11, 29 pp.
- [22] A. PAULY, *On the (semi)lattices induced by continuous reducibilities*. **Mathematical Logic Quarterly**, vol. 56 (2010), no. 5, pp. 488–502.
- [23] E. L. POST, *Recursively enumerable sets of positive integers and their decision problems*. **Bulletin of the American Mathematical Society**, vol. 50 (1944), pp. 284–316.
- [24] H. ROGERS, *Theory of Recursive Functions and Effective Computability*, McGraw-Hill, New York, 1967.
- [25] R. I. SOARE, *Turing Computability*. Theory and Applications of Computability, Springer, Berlin–Heidelberg, 2016.
- [26] R. TELGÁRSKY, *Topological games: On the 50th anniversary of the Banach–Mazur game*. **The Rocky Mountain Journal of Mathematics**, vol. 17 (1987), no. 2, pp. 227–276.
- [27] W. W. WADGE, *Reducibility and determinateness on the Baire space*. Ph.D. thesis, University of California, Berkeley, 1983.
- [28] K. WEIHRAUCH, *Type 2 recursion theory*. **Theoretical Computer Science**, vol. 38 (1985), pp. 17–33.
- [29] ———, *Computability*, EATCS Monographs on Theoretical Computer Science, vol. 9, Springer, Berlin, 1987.
- [30] ———, *Computable Analysis*, Springer, Berlin, 2000.

FACULTY OF COMPUTER SCIENCE
UNIVERSITÄT DER BUNDESWEHR MÜNCHEN
MÜNCHEN, GERMANY

and

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS
UNIVERSITY OF CAPE TOWN
CAPE TOWN, SOUTH AFRICA

E-mail: Vasco.Brattka@cca-net.de